VOL. 452 JOURNAL OF CHROMATOGRAPHY OCTOBER 28, 1988

CONTENTS

HONOUR VOLUME ON THE OCCASION OF THE 70TH BIRTHDAY OF ERICH HEFTMANN

Preface
by L. R. Snyder 1

Tswett's letter to Claparède on tropisms and taxes
by I. M. Hais (Hradec Králové, Czechoslovakia) 5

Unified theory of adsorption chromatography: gas, liquid and supercritical fluid mobile phases
by D. E. Martire (Washington, DC, U.S.A.) 17

Dispersion and selectivity indices of alkyl and alkenyl benzenes
by J. K. Haken and R. J. Smith (Kensington, N.S.W., Australia) 31

Quantitative determination of dicarboxylic acids and polyols in silicone–polyester resins
by J. K. Haken, N. Harahap and R. P. Burford (Kensington, N.S.W., Australia) 37

Analysis of short alkylsilane bonded silica gel high-performance liquid chromatographic stationary phases using hydrofluoric acid digestion and headspace analysis by capillary gas chromatography
by T. R. Floyd, N. Sagliano Jr. and R. A. Hartwick (Piscataway, NJ, U.S.A.) 43

Determination of 5-methylcytosine in DNA by gas chromatography–electron-capture detection
by D. H. Fisher and R. W. Giese (Boston, MA, U.S.A.) 51

Applications of gas chromatography to the study of terpenoid metabolism
by D. M. Satterwhite and R. B. Croteau (Pullman, WA, U.S.A.) 61

Superconductor metal oxide catalyst in a chemiluminescence chromatography detector
by E. A. McNamara, S. A. Montzka, R. M. Barkley and R. E. Sievers (Boulder, CA, U.S.A.) 75

Migration of tetrahydroisoquinoline, a possible parkinsonian neurotoxin, into monkey brain from blood as proved by gas chromatography–mass spectrometry
by T. Niwa, N. Takeda, A. Tatematsu, S. Matsuura (Nagoya, Japan), M. Yoshida (Tochigiken, Japan) and R. Nagatsu (Nagoya, Japan) 85

Identification of the more complex triacylglycerols in bovine milk fat by gas chromatography–mass spectrometry using polar capillary columns
by J. J. Myher, A. Kuksis and L. Marai (Toronto, Canada) and P. Sandra (Ghent, Belgium) 93

Studies on steroids. CCXXXVIII. Determination of bile acids in liver tissue by gas chromatography–mass spectrometry with negative ion chemical ionization detection
by J. Goto, H. Miura, M. Inada, T. Nambara, R. Nagakura and H. Suzuki (Sendai, Japan) 119

Studies of adsorption and partition effects in liquid chromatography with mixed mobile phases
by M. Borówko and M. Jaroniec (Lublin, Poland) 131

Interchangeable use of aqueous and organic solvents in a hydrophilic poly(vinyl alcohol) gel column
by S. Mori (Mie, Japan) 137

Reversed-phase high-performance liquid chromatography with a C_{18} polyacrylamide-based packing
by J. V. Dawkins and N. P. Gabbott (Loughborough, U.K.) and L. L. Lloyd, J. A. McConville and F. P. Warner (Church Stretton, U.K.) 145

Comparison of retention mechanisms of homologous series and triglycerides in non-aqueous reversed-phase liquid chromatography
by M. Martin, G. Thevenon and A. Tchapla (Palaiseau, France) 157

Expert system for chromatography
by L. Peichang and H. Hongxin (Dalian, China) 175

Considerations for using the solvent selectivity triangle approach for stationary phase characterization
by B. R. Kersten and C. F. Poole (Detroit, MI, U.S.A.) 191

Effect of mobile phase flow-rate on the recoveries and production rates in overloaded elution chromatography. A theoretical study
by S. Ghodbane and G. Guiochon (Knoxville and Oak Ridge, TN, U.S.A.) 209

Contribution of electronic effects to the lipophilicity determined by comparison of values of log *P* obtained by high-performance liquid chromatography and calculation
by J. Tipker, C. P. Groen, J. K. van den Bergh-Swart and J. H. M. van den Berg (Weesp, The Netherlands) 227

Effects of system peaks in ion-pair reversed-phase liquid chromatography for noscapine and metabolites
by M. Johansson (Solna, Sweden) and D. Westerlund (Uppsala, Sweden) 241

Measurement of the distribution coefficients of several classes of drug using reversed-phase thin-layer chromatography
by D. B. Jack, J. L. Hawker, L. Rooney, M. Beerahee, J. Lobo and P. Patel (Birmingham, U.K.) 257

Some considerations on the "charge" on a metal ion in ion-exchange chromatography
by M. Lederer (Lausanne, Switzerland) 265

Methyl green-coated column for separation of inorganic anions by ion chromatography
by R. Golombek (Stuttgart, F.R.G.) and G. Schwedt (Clausthal-Zellerfeld, F.R.G.) . . . 283

Determination of nitrite and nitrate by reversed-phase high-performance liquid chromatography using on-line post-column photolysis with ultraviolet absorbance and electrochemical detection
by M. Lookabaugh (Winchester, MA, U.S.A.) and I. S. Krull (Boston, MA, U.S.A.) . . 295

Enantioselectivity of complex formation in ligand-exchange chromatographic systems with chiral stationary and/or chiral mobile phases
by V. A. Davankov, A. A. Kurganov and T. M. Ponomareva (Moscow, U.S.S.R.) . . . 309

Determination of impurity profiles in drugs and related materials. II. Detection and quantification of diastereomeric impurity in the peptide RGH-0205 (Arg-Lys-Asp)
by S. Görög, B. Herényi, O. Nyéki, I. Schön and L. Kisfaludy (Budapest, Hungary) . . . 317

Use of hydroxypropyl- and hydroxyethyl-derivatized β-cyclodextrins for the thin-layer chromatographic separation of enantiomers and diastereomers
by D. W. Armstrong, J. R. Faulkner, Jr. and S. M. Han (Rolla, MO, U.S.A.) 323

Mixed-bed ion-exchange columns for protein high-performance liquid chromatography
by Y.-F. Maa, F. D. Antia, Z. el Rassi and Cs. Horváth (New Haven, CT, U.S.A.) . . . 331

Hydrophobic interaction of Alcian Blue with soluble and erythrocyte membrane proteins
by G. M. Ghiggeri and G. Candiano (Lavagna, Italy) and F. Ginevri (Genoa, Italy) and A. Mutti, E. Bergamaschi and R. Alinovi (Parma, Italy) and P. G. Righetti (Milan, Italy) . . 347

Purification and partial amino acid sequence of human urine protein 1. Evidence for homology with rabbit uteroglobin
by P. J. Jackson, R. Turner, J. N. Keen, R. A. Brooksbank and E. H. Cooper (Leeds, U.K.) 359

Mass spectrometry of electrophore-labeled nucleosides. Pentafluorobenzyl and cinnamoyl derivatives
by T. M. Trainor, R. W. Giese and P. Vouros (Boston, MA, U.S.A.) 369

Comparison of the chromatographic properties of sterols, select additional steroids and triterpenoids: gravity-flow column liquid chromatography, thin-layer chromatography, gas–liquid chromatography and high-performance liquid chromatography
by S. Xu, R. A. Norton, F. G. Crumley and W. D. Nes (Athens, GA, U.S.A.) 377

Quantitative microanalysis of bile acids in biological samples. Collaborative study
by F. Nakayama (Fukuoka, Japan) 399

Improved solid-phase extraction and liquid chromatography with electrochemical detection of urinary catecholamines and 5-S-L-cysteinyl-L-dopa
by T. Huang, F. Wall and P. Kabra (San Francisco, CA, U.S.A.) 409

Immunoaffinity pre-column for selective on-line sample pre-treatment in high-performance liquid chromatography determination of 19-nortestosterone
by A. Farjam, G. J. de Jong, R. W. Frei and U. A. Th. Brinkman (Amsterdam, The Netherlands) and W. Haasnoot, A. R. M. Hamers, R. Schilt and F. A. Huf (Wageningen, The Netherlands) 419

Chromatography for diagnosis of metabolic diseases
by E. Jellum (Oslo, Norway) 435

Determination of fatty acids as phenacyl esters in rat adipose tissue and blood vessel walls by high-performance liquid chromatography
by T. Hanis, M. Smrz, P. Klir, K. Macek, J. Klima, J. Base and Z. Deyl (Prague, Czechoslovakia) 443

Isolation of bile acid glucosides and N-acetylglucosaminides from human urine by ion-exchange chromatography and reversed-phase high-performance liquid chromatography
by H.-U. Marschall, G. Green, B. Egestad and J. Sjövall (Stockholm, Sweden) 459

Development of a solid-phase extraction technique for α-human atrial natriuretic peptide in human plasma
by M. A. O'Flynn, R. C. Causon, J. Brown and S. Kageyama (London, U.K.) 469

Direct enantiomeric separation of betaxolol with applications to analysis of bulk and biological samples
by A. M. Krstulović, M. H. Fouchet, J. T. Burke, G. Gillet and A. Durand (Meudon la Forêt, France) 477

High-performance liquid chromatographic analysis of the ajmalicine distribution in roots of *Catharanthus roseus* lines with different flower colours
by M. Weissenberg, A. Levy and I. Schaeffler (Bet Dagan, Israel) and E. C. Levy (Jerusalem, Israel) 485

Liquid chromatographic procedures for the analysis of compounds in the serotonergic and octoptamine pathways of lobster hemolymph
by D. A. Fadool, S. J. Cobb, G. Kass-Simon and P. R. Brown (Kingston, RI, U.S.A.) . . 491

Liquid chromatographic analysis of organic additives in copper plating baths
by S. Shohat and E. Grushka (Jerusalem, Israel) and S. Glikberg (Kiriat Malachi, Israel) . 503

High-performance liquid chromatography separations of nitrosamines. III. Conformers of N-nitrosamino acids
by H. J. Issaq, D. G. Williams, N. Schultz and J. E. Saavedra (Frederick, MD, U.S.A.) . 511

Reversed-phase C_{18} and normal-phase silica high-performance liquid chromatography of gibberellins and their methyl esters
by J.-T. Lin and A. E. Stafford (Albany, CA, U.S.A.) 519

Performance of graphitized carbon black cartridges in the extraction of some organic priority pollutants from water
by F. Mangani, G. Crescentini, P. Palma and F. Bruner (Urbino, Italy) 527

Reversed-phase liquid chromatography of astacene
by H. J. C. F. Nelis and A. P. De Leenheer (Ghent, Belgium) 535

Potential application of thin-layer chromatography and thin-layer chromatography with flame ionization detection of cholestanol in the diagnosis of cerebrotendinous xanthomatosis
by Č. Michalec and M. Ranný (Prague, Czechoslovakia) 543

Retention in sedimentation–flotation focusing field-flow fractionation using a step density gradient
by J. Jancă and N. Nováková (Brno, Czechoslovakia) 549

High-performance liquid chromatographic analysis of the alkaloid spectrum in the roots and capsules of the species and hybrids of *Papaver* section *Oxytona*
by J. Milo, A. Levy and D. Palevitch (Bet Dagan, Israel) and G. Ladizinsky (Rehovot, Israel) 563

Micellar, inclusion and metal-complex enantioselective pseudophases in high-performance electromigration methods
by J. Snopek, I. Jelínek and E. Smolková-Keulemansová (Prague, Czechoslovakia) . . . 571

Use of a double-detector system for the measurement of mobilities in zone electrophoresis
by J. L. Beckers, Th. P. E. M. Verheggen and F. M. Everaerts (Eindhoven, The Netherlands) 591

Capillary zone electrophoresis. Quantitative study of the effects of some dispersive processes on the separation efficiency
by F. Foret, M. Deml and P. Boček (Brno, Czechoslovakia) 601

Simple sampling device for capillary isotachophoresis and capillary zone electrophoresis
by Th. P. E. M. Verheggen, J. L. Beckers and F. M. Everaerts (Eindhoven, The Netherlands) 615

Separation of α-keto acids by capillary supercritical fluid chromatography as their quinoxalinol derivatives
by P. A. David and M. Novotny (Bloomington, IN, U.S.A.) 623

Author Index 631

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)